

Measurement of the Acetabular Cup Anteversion on the Simulated Radiographs

Running title: Anteversion angle of the acetabular cup

Authors: Chen-Kun Liaw^{1,2,5}, MD. PhD candidate, Rong-Sen Yang³, MD., PhD., Professor, Sheng-Mou Hou³, MD., PhD., MPH., Professor, Tai-Yin Wu⁴, MD., Chiou-Shann Fuh⁵, PhD., Professor.

Institutes: Department of Orthopaedics, Tao-Yuan General Hospital¹;

Department of Orthopaedics, En Chu Kong Hospital, Taiwan²;

Department of Orthopaedics, College of Medicine, National Taiwan University & Hospital, Taiwan³;

Taipei City Hospital, Renai Branch⁴;

Institute of Computer Science and Information Engineering, National Taiwan University⁵

Short running title: Anteversion angle of the acetabular cup

Conflict of interest statement: Each author certifies that he or she has no commercial associations (e.g., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted article.

Ethical Board Review statement: The En Chu Kong Hospital's Medical Ethics Committee has approved this study to use patients' medical records and radiographs.

Word Count (Introduction through Discussion): 1416

註解 [CORR1]: AU: Place word count here. Generally limit manuscripts to fewer than 3000 words.

Corresponding author: Correspondence to: Chiou-Shann Fuh, PhD, 4F, No. 26, Alley 2, Lane 19, Shin-Shing Street, Shan-Jia, Shu-Lin City, Taipei County, Taiwan. Phone: 886-2-23936577; Fax: 886-2-23936577; E-mail: 00901@km.eck.org.tw.
Institute of Computer Science and Information Engineering, National Taiwan University.

Chen-Kun Liaw and Rong-Sen Yang contributed equally to this work.

Abstract

The acetabulum anteversion is an important prognostic factor after THR. Widmer¹² reported a protractor to measure it on the plain radiographs. He studied the relationship between anteversion and the short axis (S) and the total length (TL) of the projected cross-section of the cup along the short axis, and approximated with linear regression. We developed our method by approximating the relationship by trigonometric mathematics. We simulated 336 radiographs with different anteversions and inclinations by our software and then measure the anteversion of the acetabular cups on these simulated radiographs by Widmer's¹² and our methods. We compared both results with the error which indicates the difference between the measured anteversion from the assumed angle on the simulated radiographs. The anteversion of the acetabular cups on the simulated radiographs ranged from 5° to 52°. The angles measured with Widmer's¹² protractor ranged from 7° to 41° (mean \pm SD =28.0° \pm 9.8°), and our methods, 5° to 51° (27.7° \pm 13.2°). The mean \pm SD of error by Widmer's¹² protractor was 5.2 \pm 2.5°, and our protractor, 0.8° \pm 0.8° (Student's t-test, p<0.0001). We also did a simple inter-observer study and found the difference between measurements of Widmer's method was less than 2°, and ours was less than 2°. The difference was smaller than the error of Widmer's method. The results showed that the error of our method was smaller than that of Widmer's¹², thus can be extended for a more precise measurement of the anteversion.

Level of Evidence: Diagnostic study, level II.

Introduction

The anteversion of acetabulum is important for function after total hip arthroplasty. Previously reported methods can be classified into three groups, the computer tomography methods^{4,9}, the trigonometric methods^{1,2,5,7,10,11}, and the protractor methods^{3,6,12}. Olivecrona et al.⁹ measured the orientation of the acetabular cups on the CT images in 10 patients. Their results showed that the anteversion angles ranged from 0° to 52° with an error of 2.9°, whereas the inclination angle ranging from 30° to 65° with an error of 1.5°.

With trigonometric method, the anteversion angles of the acetabular cups were measured using calculation equations (Appendix A). Liaw et al.⁶ applied this trigonometric method to measure the anteversion of the acetabular cups and got the mean \pm SD of error with $1.2^\circ \pm 0.57^\circ$. Additionally, Liaw et al.⁶ used his own protractor method to get the mean \pm SD of the error of $0.96^\circ \pm 0.74^\circ$. These protractor methods are more convenient than the others since they do not require a calculator or computer.

Furthermore, Liaw et al.⁶ incorporated the inverse trigonometric function into his own protractor. In practical, the most common disadvantages are to find the ends of long axis and short axis. Fabek³ applied direct measurement using a protractor that was designed without any incorporation of trigonometric function. However, the examiner usually has difficulty in following the long arc of the circles during the measurement. Widmer¹² invented his own protractor through his linear regression equation. The user can apply for direct measurement without the need of finding the ends of the long axis first. Widmer¹² mentioned that the only disadvantage is its imprecision that was due to oblique radiographic projection on various acetabulum abduction angles and the adoption

of a linear regression equation. He did not recommend the usage of his own protractor if highly precise measurements are needed. This raised some questions. What is the source of the error? Can we improve it?

The study aims to investigate the relationship curve mathematically and to eliminate the error caused by oblique projection. The measured angles and the precision error will be compared with those of the Widmer's¹² results.

Materials and Methods

At the given distance of 105 cm from x-ray tubes to subjects, Widmer¹² found a relationship between anteversion and the short axis (S) and the total length (TL) of the projected cross-section of the cup along the short axis by linear regression.

$$\text{Anteversion} = 48.5 \cdot (S/TL) - 0.3$$

In our methods, we investigated the mathematical relationship between radiographic version β and S/TL-ratio is shown in Equation (1). The detailed deduction process was shown in Appendix A.

$$\beta = \sin^{-1}(S / l) = \sin^{-1}((S/TL\text{-ratio}) / (2 - (S/TL\text{-ratio}))) \quad (1)$$

To eliminate the error caused by oblique projection, we applied the Equation (2). The detailed deduction process was shown in Appendix B.

$$\beta = \tan^{-1}(\tan(\tan^{-1}(\tan(\sin^{-1}((S/TL\text{-ratio}) / (2 - (S/TL\text{-ratio})))))) \csc \gamma) + 5.46^\circ \sin \gamma \quad (2)$$

Through Equations 2 we reproduced Widmer's¹² results that were shown in Fig. 1 and Table 1. The results were quite the same as the data shown by Widmer¹².

We further used the mathematic model to calculate the error of Widmer's¹² linear regression equation (Fig. 2), and improved the precision by the following two methods.

First, we applied the protractor on the hip-centered radiographs that eliminated the error caused by oblique projection. If we used the radiograph centered on the symphysis pubis for measurement, we corrected by Equation 2.

Second, we improved the precision by a mathematic model. Widmer's¹² method used linear regression method to approximate the curve. The precision was good in linear region of the whole curve, but bad in the non-linear region. The mathematic model fully approximated the curve, thus improved the precision.

Base on these two points, we developed our protractor through Equation 1 (Fig. 3A).

In order to determine the accuracy, we made a Widmer's¹² protractor through his linear regression equation ($y = 48.05x - 0.3$) and our protractor (Fig. 3B). We simulated 336 total hip arthroplasty radiographs with 48 different anteversions ranging from 5°–52° and seven different inclinations (30°, 35°, 40°, 45°, 50°, 55°, 60°) using our simulation program. We removed the femoral heads and necks in our simulated radiographs to eliminate the occluding effects. We used these two protractors to measure anteversions on these simulated radiographs. We found first the perpendicular bisector of the long axis of the acetabular cup. Then we found three intersection points between the perpendicular bisector and the ellipse by the rim of the acetabular cup or the hemisphere curve by outer shell. Then we applied the protractors to read the anteversion angle (Fig. 3C & 3D). Widmer's¹² protractor had a built-in correction of the projection obliquity. For comparison, we corrected the anteversion centered at hip to anteversion centered at symphysis pubis by following procedure. First we converted the real anteversion to anatomic anteversion, subtracting 5.46°, and then converting back to radiographic

anteversion. The anteversion angles on the simulated radiographs were measured by one author in a random order using either method. The precision error was calculated from the difference between the measured angles and the assumed angles of these simulated radiographs. These results were compared by Student's t-test.

To justify our improvement, we did an inter-observer difference study by randomly selecting 10 hip arthroplasty radiographs and measured the radiographic anteversion with our method and Widmer's method each twice by two of the authors. Then we calculated absolute difference of two measurements. If the difference was larger than the error of Widmer's error, our improvement made little sense.

Results

The angles measured with Widmer's¹² method ranged from 7° to 41° (mean ± SD =28.0° ± 9.8°), and for our methods, 5° to 51° (27.7 ±13.2°). After oblique projection correction, the real radiographic anteversion (centered at symphysis pubis) used for Widmer's method ranged from 0.3° to 49.0°. The error of Widmer's protractor ranged from 0° to 8.7°, and the mean ± SD is 5.2 ± 2.5°(Fig. 4A); the range with our protractor, 0° to 3°, and mean ± SD, 0.8° ± 0.8°(Fig. 4B)(Student's t-test, p<0.0001).

For the inter-observer study, the radiographic anteversion measured by Widmer's method twice ranged from 3° to 21° (mean ± SD =12.3° ± 5.9°), and by ours twice, 2° to 16° (8.7° ± 4.7°). The absolute difference between two measurements of Widmer's method ranged from 0° to 2° (mean ± SD =0.5° ± 0.7°), and of ours, 0° to 1° (0.5° ± 0.7°).

Discussion

Measuring anteversion is a cumbersome work for a medical doctor. In our experience, Widmer¹² designed a rather convenient method as compared with others

whereas his method incorporated a potential imprecision. Therefore, to improve the imprecision of his method may refine the measurement.

With application of perpendicular bisector for the measurement and mathematical equations, our modified protractor has significantly reduced the error by using our own protractor for the measurement of the anteversion of the acetabular cups. The improvement was statistically significant. The error of Widmer's¹² method was mainly related to inclination angle and anteversion angle. The correlation between error and inclination was caused by that Widmer ignored the influence of inclination when correcting oblique projection. The correlation between error and anteversion was because that Widmer used linear regression to approximate the curve. This finding in this study correlated well with his previous report. Our method improved the precision in both types of error. However, our method has larger error when anteversion increased. The reason was we underestimated the short axis (S). When anteversion increased, the outer edge became blurred. If we measured with the inner edge, thus we underestimated the short axis (S). Fortunately this error was small in our study, only 3° when anteversion larger than 45°. The intra-observer difference of Widmer's method was between 0° to 2°, and of ours 0° to 2°, which was smaller than the error of Widmer's method. Our improvement did make difference in this situation.

The range of the simulated radiographs' anteversion is between 5° to 51° for our method and 0.3° to 49.0° for Widmer's method. In study of Olivecrona et al⁹, the range of anteversion is between 0° to 52° and inclination is between 30° to 65°.² Therefore we chose the aforementioned range of anteversion for measurement in this study.

Since we had to face the possible error caused by the projection, the limitation of this study was that we need a basic assumption of the perfect hemi-ball shape for the acetabulum. If not, our method was not suitable. In that situation, Liaw's⁶ and Fabeck's³ protractors were preferred. Otherwise, our improvement had significantly reduced the error, thus can be used in precise measurement of the anteversion.

References

1. Ackland MK, Bourne WB, Uthoff HK. Anteversion of the acetabular cup: measurement of angle after total hip replacement. *J Bone Joint Surg Br.* 1986;68:409-413.
2. Coventry MB, Beckenbaugh RD, Nolan DR, Ilstrup DM. 2,012 total hip arthroplasties: a study of postoperative course and early complications. *J Bone Joint Surg Am.* 1974;56:273-284.
3. Fabeck L, Farrokh D, Tolley M, Descamps PY, Gebhart M, Delince P. A method to measure acetabulum cup anteversion after total hip replacement. *Acta Orthop Belg.* 1999;65:485-491.
4. Jaramaz B, DiGioia AM, Blackwell M, Nolan DR. Computer assisted measurement of cup placement in total hip replacement. *Clin Orthop* 354:70–80, 1998.
5. Lewinnek GE, Lewis JL, Tarr R, Compere CL, Zimmerman JR. Dislocations after total hip-replacement arthroplasties. *J Bone Joint Surg Am.* 1978;60:217-220.
6. Liaw CK, Hou SM, Yang RS, et al. A New Tool for Measuring Cup Orientation in Total Hip Arthroplasties from Plain Radiographs. *Clin Orthop.* 2006;451:134–139.
7. McLaren RH. Prosthetic Hip Angulations. *Radiology.* 1973;107:705-706.
8. Murray DW. The definition and measurement of acetabular orientation. *J Bone Joint Surg Br.* 1993;75:228-232.
9. Olivecrona H, Weidenhielm L, Olivecrona L, Beckman MO, et al. A new CT method for measuring cup orientation after total hip arthroplasty: A study of 10 patients. *Acta Orthop Scand.* 75:252–260, 2004.

10. Pradhan R. Planar anteversion of the acetabular cup as determined from plain antero-posterior radiographs. *J Bone Joint Surg Br.* 1999;81:431-435.
11. Visser JD, Konings JG. A new method for measuring angles after total hip arthroplasty: a study of the acetabular cup and femoral component. *J Bone Joint Surg Br.* 1981;63:556-559.
12. Widmer KH. A simplified method to determine acetabular cup anteversion from plain radiographs. *J Arthroplasty.* 2004;19:387-390.

Table

Table 1. By Equations 4 and 7, the relationship between S/TL ratio and anteversion is shown. This results are similar to Widmer's¹² report.

Legends

Fig. 1 The relationship between S/TL ratio and radiographic anteversion. This figure is derived from Equations 4 and 7, and the inclination angle equals 45°.

Fig. 2. Estimated error of Widmer's¹² linear regression equation. With our mathematic model, we calculate the ideal anteversion from S/TL ratio. The error is the difference between the ideal anteversion and Widmer's linear regression anteversion

Fig. 3A. Our protractor developed through Equation 1.

Fig. 3B. Widmer's¹² protractor made according to his linear regression equation ($y = 48.05x - 0.3$).

Fig. 3C. The simulated radiographs are printed on papers. Then we use our protractor to measure the radiographic anteversion.

Fig. 3D. The simulated radiographs are printed on papers. Then we use Widmer's¹² protractor to measure the radiographic anteversion.

Fig. 4A. The error of Widmer's¹² method. Clearly, the error is related to inclination angle and anteversion angle.

Fig. 4B. The error of our method. The error is slightly related to anteversion angle.